


# Can BOTOX® Really Improve Your Overactive Bladder Symptoms?



## The Results Speak for Themselves


Patients receiving BOTOX® reported a significant reduction in the number of their daily leakage episodes.



At week 12, many of these patients saw their daily leakage episodes reduced by half or more following BOTOX® treatment.\*



The majority of patients receiving BOTOX® had at least **50%** reduction of daily leakage episodes.



Most of these patients reduced their leakage episodes by **75%**.



**APPROXIMATELY 1 IN 4** of them reported they were experiencing no leakage episodes.

In clinical trials, one BOTOX® treatment provided symptom improvement for up to 6 months. Your doctor will work with you to develop a re-treatment plan.

\* Reduction of Urinary Incontinence (UI) episodes in:

### Study 1: BOTOX® n = 280; placebo n = 277;

- Patients with U50% reduction: BOTOX® 57.5%, placebo 28.9% (P < .001)
- Patients with U75% reduction: BOTOX® 44.6%, placebo 15.2% (P < .001)
- Patients with 100% reduction: BOTOX® 22.9%, placebo 6.5% (P < .001)

### Study 2: BOTOX® n = 277; placebo n = 27

- Patients with U50% reduction: BOTOX® 63.5%, placebo 33.2% (P < .001)
- Patients with U75% reduction: BOTOX® 47.3%, placebo 20.3% (P < .001)
- Patients with 100% reduction: BOTOX® 31.4%, placebo 10.3% (P < .001)

### Study 3: BOTOX® n = 145; placebo n = 60

- Patients with U50% reduction: BOTOX® 77.2%, placebo 33.3% (P < .001)
- Patients with U75% reduction: BOTOX® 55.2%, placebo 21.7% (P < .001)
- Patients with 100% reduction: BOTOX® 33.8%, placebo 11.7% (P < .001)

Over **50%** of patients have a **75%** or greater reduction in urinary incontinence episodes

By significantly reducing daily leakage episodes, BOTOX® treatment may make a positive impact on your daily life.

Ask your doctor if BOTOX treatment is right for you.